

XCS Service v1.00
con .cas
9351006 COM

Welcome to STN International! Enter x:x

LOGINID:ssspta1811bxp

PASSWORD:

khbed465

* * * * * RECONNECTED TO STN INTERNATIONAL * * * * *
SESSION RESUMED IN FILE 'REGISTRY' AT 11:10:36 ON 09 JAN 97
FILE 'REGISTRY' ENTERED AT 11:10:36 ON 09 JAN 97
COPYRIGHT (C) 1997 American Chemical Society (ACS)

=> d his

(FILE 'HOME' ENTERED AT 10:14:44 ON 09 JAN 97)

FILE 'REGISTRY' ENTERED AT 10:15:04 ON 09 JAN 97

E CETRORELIX
L1 1 S E3
E BOMBESIN
L2 242 S E3
E BOMBESIN/CN
L3 1 S E3
E PROTIRELIN
L4 1 S E3

=> e somatostatin/cn

E1 1 SOMATORELIN/CN
E2 1 SOMATOSALM/CN
E3 2 --> SOMATOSTATIN/CN
E4 1 SOMATOSTATIN (AMIA CALVA REDUCED)/CN
E5 1 SOMATOSTATIN (AMPHIUMA TRIDACTYLUM)/CN
E6 1 SOMATOSTATIN (ANGUILLA ANGUILLA)/CN
E7 1 SOMATOSTATIN (HUMAN)/CN
E8 1 SOMATOSTATIN (HYDROLAGUS COLLIEI)/CN
E9 1 SOMATOSTATIN (LOPHIUS AMERICANUS REDUCED)/CN
E10 1 SOMATOSTATIN (LOPHIUS AMERICANUS)/CN
E11 1 SOMATOSTATIN (MAMMALIAN-REDUCED)/CN
E12 1 SOMATOSTATIN (MYXINE GLUTINOSA)/CN

=> s e3

L5 2 SOMATOSTATIN/CN

=> s l1 or l3 or l4 or l5

L6 5 L1 OR L3 OR L4 OR L5

=> file hcaplus

FILE 'HCAPLUS' ENTERED AT 11:12:18 ON 09 JAN 97
USE IS SUBJECT TO THE TERMS OF YOUR CUSTOMER AGREEMENT
COPYRIGHT (C) 1997 AMERICAN CHEMICAL SOCIETY (ACS)

FILE COVERS 1967 - 9 Jan 1997 VOL 126 ISS 2
FILE LAST UPDATED: 9 Jan 1997 (970109/ED)

For information on changes to CA General Subject Index
terms in 1997, see the CAS Web search aid at

STN 08/468145; 1/9/97

<http://www.cas.org/terms/vocab.html>

=> s l6

L7 17904 L6

=> s l7 and lyophil?

11647 LYOPHIL?

L8 35 L7 AND LYOPHIL?

=> s l8 and py<1994

11518711 PY<1994

L9 30 L8 AND PY<1994

=> s l9 and water

1144296 WATER

L10 4 L9 AND WATER

=> d bib ab 1-

L10 ANSWER 1 OF 4 HCAPLUS COPYRIGHT 1997 ACS

AN 1992:113531 HCAPLUS

DN 116:113531

TI Solvent systems for production of drug microspheres

IN Yamakawa, Ichiro; Machida, Ryoichi; Watanabe, Sumio

PA Eisai Co., Ltd., Japan

SO Eur. Pat. Appl., 12 pp.

CODEN: EPXXDW

PI EP 461630 A2 911218

DS R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE

AI EP 91-109610 910612

PRAI JP 90-152849 900613

JP 90-152850 900613

JP 91-48579 910222

DT Patent

LA English

AB Upon incorporation of a drug into microspheres by oil/**water** solvent evapn., a mixed solvent of .gtoreq.1 **water**-insol. solvent and .gtoreq.1 **water**-miscible solvent is used as a solvent of an oil phase. The oil phase may also comprise fatty acids or salts thereof, glycerin fatty acid esters, and propylene glycol fatty acid esters. The resulting microspheres contain the drug at a high concn. and slowly release from the initial stage after administration. Thus, neurotensin analog 20 and poly(DL-lactic acid) 200 mg were dissolved in 0.6 mL of methylene chloride/EtOH (5:1 by vol.) and the resulting soln. was dispersed in 200 mL of a 0.5 % aq. soln. of PVA. The dispersion was stirred for 3 h to conduct oil/**water** solvent evapn. so that an oil phase was solidified. Microspheres thus formed were collected by a centrifugal separator and **lyophilized** into powder. The drug content in the microspheres was 21.5 %, vs. 10.4 % for the control microspheres produced in a similar manner except that 0.6 mL of methylene chloride was used instead of the mixed solvent.

L10 ANSWER 2 OF 4 HCAPLUS COPYRIGHT 1997 ACS

AN 1990:145467 HCAPLUS

DN 112:145467

TI Preparation of a somatostatin injection in hospital pharmacy

AU Hoshino, Nobuo; Yamakawa, Masayuki; Nishida, Yumiko; Ono, Takeshi; Shimakawa, Harumi
CS Hosp. Pharm., Shiga Univ. Med. Sci., Otsu, 520-21, Japan
SO Yakuzai-gaku (1989), 49(3), 263-7
CODEN: YAKUA2; ISSN: 0372-7629
DT Journal
LA Japanese
AB Two types of somatostatin (I) injections, I soln. and lyophilized I, were prepd. and investigated from the view points of both stability in moist heat sterilization and stability during storage under various environmental conditions. In I soln., I was dissolved in 50 mM acetate buffer (pH 4 and 5) and phosphate buffer (pH 6, 7 and 8) at a concn. of 250 μ g/mL, dispensed into ampuls and then sterilized by moist heat. In lyophilized I, I was dissolved in water, dispensed into ampuls, and then lyophilized. In order to measure the stability of the preps., residual somatostatin content was detd. by HPLC and residual biol. potency by bioassay. In moist heat sterilization, I soln. was stable at pH 4 but not at higher pH. During 1 yr storage, I soln. at 3.degree. in the dark was more stable than at room temp. (20.degree.), and its decompn. was <5% under any of the three conditions (at 3.degree. in the dark, at room temp. in the dark and at room temp. under scattered light) for 1 yr.

L10 ANSWER 3 OF 4 HCAPLUS COPYRIGHT 1997 ACS
AN 1984:17886 HCAPLUS
DN 100:17886
TI Characterization of rat placental TRH-like material and the ontogeny of placental and fetal brain TRH
AU Shambaugh, George E., III; Kubek, Michael; Wilber, John F.
CS Sch. Med., Northwestern Univ., Chicago, IL, 60611, USA
SO Placenta (1983), 4(4), 329-37
CODEN: PLACDF; ISSN: 0143-4004
DT Journal
LA English
AB TRH [24305-27-9] immunol. and biol. activity was characterized in rat placental exts. Placentas were extd. sequentially with 2N AcOH and glacial AcOH. The lyophilized residues were further extd. with 90% MeOH, and the exts. were then dried and reextd. with distd. water. When a 100,000-g supernatant fraction of this ext. was utilized for characterization, TRH extd. from the placenta was similar to synthetic TRH by 4 criteria: parallelism of immunoassay inhibition curves, similarity of elution vols. after Sephadex G-10 chromatog., TRH biol. activity (TSH release from rat pituitaries in vitro), and identity on high-pressure liq. chromatog. Between the 16th and 20th day of gestation, placental TRH activity increased nearly 3-fold, from 10.9 to 29.7 pg/mg protein. Changes of a similar magnitude were apparent in the fetal brain (6.0 to 18.9 pg/mg protein). Hence, TRH activity in the rat placenta increases with gestational age in a pattern similar to that described previously for certain placental protein hormones and developmental changes in fetal brain TRH.

L10 ANSWER 4 OF 4 HCAPLUS COPYRIGHT 1997 ACS
AN 1981:154861 HCAPLUS
DN 94:154861
TI Production of antigens containing bradykinin and somatostatin

PA Otsuka Pharmaceutical Co., Ltd., Japan
SO Jpn. Kokai Tokkyo Koho, 7 pp.
CODEN: JKXXAF
PI JP 56016414 810217 Showa
AI JP 79-92781 790720
DT Patent
LA Japanese
AB An Ascaris suum ext. is a more effective carrier of haptens like somatostatin and bradykinin than serum albumins. Thus, an antigen, somatostatin-Ascaris ext.-carbodiimide complex, was prepd. by reacting somatostatin (1.0 mg) dissolved in 200 .mu.L water with Ascaris ext. (10.3 mg) in a .5 mM borate buffer (0.15M NaCl, pH 8.0), then adding 50 .mu.g carbodiimide in 500 .mu.L water, and adjusting the pH at 6.5 with 1N HCl. The reaction mixt. was dialyzed at 4.degree. for 5 h against water; this dialysis was repeated 4 times. The yield of the antigen complex after the lyophilization was 12.4 mg.

=> d his

(FILE 'HOME' ENTERED AT 10:14:44 ON 09 JAN 97)

FILE 'REGISTRY' ENTERED AT 10:15:04 ON 09 JAN 97

E CETRORELIX
L1 1 S E3
E BOMBESIN
L2 242 S E3
E BOMBESIN/CN
L3 1 S E3
E PROTIRELIN
L4 1 S E3
E SOMATOSTATIN/CN
L5 2 S E3
L6 5 S L1 OR L3 OR L4 OR L5

FILE 'HCAPLUS' ENTERED AT 11:12:18 ON 09 JAN 97

L7 17904 S L6
L8 35 S L7 AND LYOPHIL?
L9 30 S L8 AND PY<1994
L10 4 S L9 AND WATER

=> file reg

FILE 'REGISTRY' ENTERED AT 11:16:36 ON 09 JAN 97

USE IS SUBJECT TO THE TERMS OF YOUR CUSTOMER AGREEMENT
COPYRIGHT (C) 1997 American Chemical Society (ACS)

STRUCTURE FILE UPDATES: 3 JAN 97 HIGHEST RN 184695-79-2
DICTIONARY FILE UPDATES: 8 JAN 97 HIGHEST RN 184695-79-2

TSCA INFORMATION NOW CURRENT THROUGH JUNE 1996

Please note that search-term pricing does apply when
conducting SmartSELECT searches.

=> s acetic acid/cn

L11 1 ACETIC ACID/CN

=> s l8 and l11
1 LYOPHIL?
L12 0 L8 AND L11

=> file caplus
FILE 'CAPLUS' ENTERED AT 11:17:24 ON 09 JAN 97
USE IS SUBJECT TO THE TERMS OF YOUR CUSTOMER AGREEMENT
COPYRIGHT (C) 1997 AMERICAN CHEMICAL SOCIETY (ACS)

FILE COVERS 1967 - 9 Jan 1997 VOL 126 ISS 2
FILE LAST UPDATED: 9 Jan 1997 (970109/ED)

To help control your online searching costs, consider using the
HCAPLUS file when using the FSEARCH command or when conducting
SmartSELECT searches with large numbers of terms.

For information on changes to CA General Subject Index
terms in 1997, see the CAS Web search aid at
<http://www.cas.org/terms/vocab.html>

=> s l8 and l11
17904 L6
11647 LYOPHIL?
45618 L11
L13 1 L8 AND L11

=> d bib ab

L13 ANSWER 1 OF 1 CAPLUS COPYRIGHT 1997 ACS
AN 1994:587330 CAPLUS
DN 121:187330
TI Preparation of a cetorelix **lyophilized** composition
IN Engel, Juergen; Sauerbier, Dieter; Wichert, Burkhard; Reissmann,
Thomas
PA Asta Medica AG, Germany
SO Eur. Pat. Appl., 5 pp.
CODEN: EPXXDW
PI EP 611572 A2 940824
DS R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LI, LU, MC, NL, PT,
SE
AI EP 94-101672 940204
PRAI DE 93-4305225 930219
DT Patent
LA German
AB A **lyophilizate** of a peptide with 3-15 amino acid residues
(e.g. cetorelix) and .gtoreq.1 optional matrix materials (e.g.
mannitol) is prepd. by dissolving in 100-10,000 wt. parts AcOH,
dilg. with water, and **lyophilizing** the resulting soln.
The **lyophilizate** is useful for prepn. of a medication for
treatment of female infertility and protection of the gonads from
the follicular hyperstimulation seen with other infertility
treatments.

=> d bib ab l10 1-
NO ANSWERS DISPLAYED.
THE ANSWER SET WAS CREATED IN FILE 'HCAPLUS'.
USE THE FILE COMMAND TO CHANGE TO THE CORRECT FILE.

You have entered a file that is not in the current file environment.
Enter "DISPLAY HISTORY" to see a list of the files in the current environment.

=> filoe hcaplus

'FILOE' IS NOT A RECOGNIZED COMMAND

The previous command name entered was not recognized by the system.

For a list of commands available to you in the current file, enter

"HELP COMMANDS" at an arrow prompt (=>).

=> file hcaplus

FILE 'HCAPLUS' ENTERED AT 11:19:47 ON 09 JAN 97

USE IS SUBJECT TO THE TERMS OF YOUR CUSTOMER AGREEMENT

COPYRIGHT (C) 1997 AMERICAN CHEMICAL SOCIETY (ACS)

FILE COVERS 1967 - 9 Jan 1997 VOL 126 ISS 2

FILE LAST UPDATED: 9 Jan 1997 (970109/ED)

For information on changes to CA General Subject Index
terms in 1997, see the CAS Web search aid at
<http://www.cas.org/terms/vocab.html>

=> d bib ab 110 1-

L10 ANSWER 1 OF 4 HCAPLUS COPYRIGHT 1997 ACS

AN 1992:113531 HCAPLUS

DN 116:113531

TI Solvent systems for production of drug microspheres

IN Yamakawa, Ichiro; Machida, Ryoichi; Watanabe, Sumio

PA Eisai Co., Ltd., Japan

SO Eur. Pat. Appl., 12 pp.

CODEN: EPXXDW

PI EP 461630 A2 911218

DS R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE

AI EP 91-109610 910612

PRAI JP 90-152849 900613

JP 90-152850 900613

JP 91-48579 910222

DT Patent

LA English

AB Upon incorporation of a drug into microspheres by oil/water solvent evapn., a mixed solvent of .gtoreq.1 water-insol. solvent and .gtoreq.1 water-miscible solvent is used as a solvent of an oil phase. The oil phase may also comprise fatty acids or salts thereof, glycerin fatty acid esters, and propylene glycol fatty acid esters. The resulting microspheres contain the drug at a high concn. and slowly release from the initial stage after administration. Thus, neurotensin analog 20 and poly(DL-lactic acid) 200 mg were dissolved in 0.6 mL of methylene chloride/EtOH (5:1 by vol.) and the resulting soln. was dispersed in 200 mL of a 0.5 % aq. soln. of PVA. The dispersion was stirred for 3 h to conduct oil/water solvent evapn. so that an oil phase was solidified. Microspheres thus formed were collected by a centrifugal separator and lyophilized into powder. The drug content in the microspheres was 21.5 %, vs. 10.4 % for the control microspheres produced in a similar manner except that 0.6 mL of methylene chloride was used instead of the mixed solvent.

L10 ANSWER 2 OF 4 HCAPLUS COPYRIGHT 1997 ACS
AN 1990:145467 HCAPLUS
DN 112:145467
TI Preparation of a somatostatin injection in hospital pharmacy
AU Hoshino, Nobuo; Yamakawa, Masayuki; Nishida, Yumiko; Ono, Takeshi; Shimakawa, Harumi
CS Hosp. Pharm., Shiga Univ. Med. Sci., Otsu, 520-21, Japan
SO Yakuzaigaku (1989), 49(3), 263-7
CODEN: YAKUA2; ISSN: 0372-7629
DT Journal
LA Japanese
AB Two types of somatostatin (I) injections, I soln. and lyophilized I, were prepd. and investigated from the view points of both stability in moist heat sterilization and stability during storage under various environmental conditions. In I soln., I was dissolved in 50 mM acetate buffer (pH 4 and 5) and phosphate buffer (pH 6, 7 and 8) at a concn. of 250 .mu.g/mL, dispensed into ampuls and then sterilized by moist heat. In lyophilized I, I was dissolved in water, dispensed into ampuls, and then lyophilized. In order to measure the stability of the preps., residual somatostatin content was detd. by HPLC and residual biol. potency by bioassay. In moist heat sterilization, I soln. was stable at pH 4 but not at higher pH. During 1 yr storage, I soln. at 3.degree. in the dark was more stable than at room temp. (20.degree.), and its decompn. was <5% under any of the three conditions (at 3.degree. in the dark, at room temp. in the dark and at room temp. under scattered light) for 1 yr.

L10 ANSWER 3 OF 4 HCAPLUS COPYRIGHT 1997 ACS
AN 1984:17886 HCAPLUS
DN 100:17886
TI Characterization of rat placental TRH-like material and the ontogeny of placental and fetal brain TRH
AU Shambaugh, George E., III; Kubek, Michael; Wilber, John F.
CS Sch. Med., Northwestern Univ., Chicago, IL, 60611, USA
SO Placenta (1983), 4(4), 329-37
CODEN: PLACDF; ISSN: 0143-4004
DT Journal
LA English
AB TRH [24305-27-9] immunol. and biol. activity was characterized in rat placental exts. Placentas were extd. sequentially with 2N AcOH and glacial AcOH. The lyophilized residues were further extd. with 90% MeOH, and the exts. were then dried and reextd. with distd. water. When a 100,000-g supernatant fraction of this ext. was utilized for characterization, TRH extd. from the placenta was similar to synthetic TRH by 4 criteria: parallelism of immunoassay inhibition curves, similarity of elution vols. after Sephadex G-10 chromatog., TRH biol. activity (TSH release from rat pituitaries in vitro), and identity on high-pressure liq. chromatog. Between the 16th and 20th day of gestation, placental TRH activity increased nearly 3-fold, from 10.9 to 29.7 pg/mg protein. Changes of a similar magnitude were apparent in the fetal brain (6.0 to 18.9 pg/mg protein). Hence, TRH activity in the rat placenta increases with gestational age in a pattern similar to that described previously for certain placental protein hormones and developmental changes in fetal brain TRH.

L10 ANSWER 4 OF 4 HCAPLUS COPYRIGHT 1997 ACS
AN 1981:154861 HCAPLUS
DN 94:154861
TI Production of antigens containing bradykinin and somatostatin
PA Otsuka Pharmaceutical Co., Ltd., Japan
SO Jpn. Kokai Tokkyo Koho, 7 pp.
CODEN: JKXXAF
PI JP 56016414 810217 Showa
AI JP 79-92781 790720
DT Patent
LA Japanese
AB An Ascaris suum ext. is a more effective carrier of haptens like somatostatin and bradykinin than serum albumins. Thus, an antigen, somatostatin-Ascaris ext.-carbodiimide complex, was prepd. by reacting somatostatin (1.0 mg) dissolved in 200 .mu.L water with Ascaris ext. (10.3 mg) in a 5 mM borate buffer (0.15M NaCl, pH 8.0), then adding 50 .mu.g carbodiimide in 500 .mu.L water, and adjusting the pH at 6.5 with 1N HCl. The reaction mixt. was dialyzed at 4.degree. for 5 h against water; this dialysis was repeated 4 times. The yield of the antigen complex after the lyophilization was 12.4 mg.

=> log y